

Domande tecniche prova orale

1

- TRAUMI DISTORSIVI DELLA CAVIGLIA: DECORSO E TRATTAMENTO
- POSIZIONAMENTO A LETTO NEI VARI DECUBITI DI UN PAZIENTE POST-ICTUS IN FASE ACUTA: SIGNIFICATO DEL PROGRAMMA RIABILITATIVO

2

- IL RAZIONALE DEL TRATTAMENTO POST CHIRURGICO DELLA PROTESI DI TOTALE DEL GINOCCHIO NEL PAZIENTE ANZIANO
- LA RIABILITAZIONE DELL'EMIPLEGICO ADULTO: METODICHE RIABILITATIVE

Roberto Scarpelli

3

- LE COMPLICANZE PIU FREQUENTI NELLA SPALLA PLEGICA: PREVENZIONE E TRATTAMENTO
- IL RAZIONALE DEL TRATTAMENTO POST CHIRURGICO NELLE FRATTURE DI FEMORE NELL'ANZIANO

Stefano Teta

Al. 8

Al. 8

Al. 8

Prova Informatica

Paolo

N1- Che estensione hanno i file di word?

N2- Come si crea una cartella?

Cartella

N3- Cosa significa WWW?

WWW

W

World Wide Web

World Wide Web

Prova lingua inglese

1

Carly

Treatment options include angioplasty, bypass or amputation of the limb, when life or limb is threatened. People with CLI have a high risk of mortality and morbidity. The mortality rates during a surgical admission are approximately 5%. Within one year of surgery, the mortality rate rises to 22%. Postoperative complications are as high as 30% and readmission rates vary between 7% to 18% in people with CLI.

2

The use of prehabilitation is gaining momentum, particularly in elderly patients undergoing surgery and patients undergoing colorectal cancer surgery, as a means of optimising fitness to improve the prognosis for people undergoing the physiological stress of surgery. People with PAD are characterised by poor mobility and physical function and have a lower level of fitness as a result of disease progression.

3

John to the

This review aimed to compare prehabilitation with usual care (defined as a preoperative assessment, including blood and urine tests). The key outcomes were postoperative complications, mortality and readmissions within 30 days of the surgical procedure, and one-year survival rates.

[Signature]

Paul John O'Neil

[Signature]